

Review

Prevalence of Pediculosis Capitis in Indonesia

Reggi First Trasia 1

¹ Parasitology Division, Faculty of Medicine, Universitas Sultan Ageng Tirtayasa

Corresponding author: reqgi.first@untirta.ac.id

ABSTRACT

Head lice infestation is still a health problem in Indonesia. Currently, there has been a lot of research on the incidence of pediculosis capitis. However, there are still few articles that systematically review the incidence of head lice infestation. The purpose of writing this article is to examine cases of pediculosis capitis in several regions in Indonesia in order to obtain systematic and comprehensive prevalence data. A literature search was carried out in October 2021 through Google Scholar and obtained 13 relevant articles. Based on the above review, it can be concluded that the highest incidence of pediculosis capitis is in Islamic boarding schools, attacking elementary school-aged children, with a prevalence range of 29.3%-88.9% and can occur throughout Indonesia. This condition can be caused by a lack of personal hygiene, so that it can continue to be a source of infestation for children around it.

Keywords: Prevalence, pediculosis capitis, parasitic infestation, Indonesia, tropical disease.

INTRODUCTION

Head lice (*Pediculus humanus capitis*) are insects that live in between the hair and attached to the human scalp. These animals are very small in size and survive by sucking blood through the human scalp, and reproduce by laying eggs and nesting their eggs on strands of hair, especially at the base of the hair. The impact of *Pediculus humanus capitis* on health is that it can cause scabs on the head and Pediculosis capitis. In children, there are health problems, namely Pediculosis capitis which can cause anemia which makes children lethargic, sleepy in class, affects learning performance and reduces children's confidence.¹

The Centers for Disease Control and Prevention (CDC) in 2016 stated that *Pediculus humanus capitis* infestation spread throughout the world with the highest incidence in children aged 3 to 11 years. *Pediculus humanus capitis* can live in human hair due to several factors, namely the lack of personal hygiene, especially the cleanliness of the hair and scalp, and can also be infected through daily activities between other individuals who have been infected with *Pediculus humanus capitis*.²

In the United States, it is reported that about 6 million to 12 million children are affected by Pediculosis capitis, especially in girls aged 3 to 11 years. The very high infection rate in 2015 was reported to be more than 70% in Pakistan. In several regions in Indonesia, many studies have been conducted on the incidence of pediculosis capitis. However, there are still few articles that systematically review the incidence of head lice infestation. The purpose of writing this article is to examine cases of pediculosis capitis in several regions in Indonesia in order to obtain systematic and comprehensive prevalence data..

METHODS

The literature search was carried out in October 2021 through Google Scholar. The search keywords were "pediculosis and prevalence". The inclusion criteria for articles were information on the prevalence of pediculosis, research conducted in Indonesia, and articles published between 2020-2021. The exclusion criteria were that there were duplications and they were not fully accessible. The publications selected in the selection of titles and abstracts were extracted using a standard format table and processed using a Microsoft Excel spreadsheet. The extracted data are the author, year of

publication, journal, research location and conclusions. The results are then presented qualitatively.

RESULTS

From the results of a search of 135 articles, there were 13 articles that met the inclusion and exclusion criteria. A critical study was conducted on selected articles published by accredited national journals. Analysis of the prevalence of pediculosis in various regions in Indonesia in 2020-2021 can be seen in table 1.

Table 1. Prevalence of Pediculosis in Indonesia 2020-2021

Research	Prevalen ce	Location	Reference
Prevalence of pediculosis kapitis and risk factors for infestation in children in elementary school No. 6 Darmasaba.	59.7% in elementa ry school subjects	Kec. Abiansem al, Kab. Badung, Bali	(Suweta NPTB, et al. 2021)
Relationship of risk factors with the proportion of pediculus humanus capitis infection in Muhammadi yah 1 elementary school students in Medan City	29.3% on elementa ry school subjects	Kota Medan, Sumatera Utara	(Syarbaini S, et al. 2021)
Description of personal hygiene behavior related to head lice infestation (pediculus humanus capitis) at Pondok Pesantren	88.2% on the subject of santriwat i	Kab. Kampar, Riau	(Analdi V, et al. 2021)

Anshor Al-			
Sunnah			
Riau			
The	66% on	Kab.	(Sulistyaningt
relationship	santriwat	Rembang,	yas, et al.
between	i subjects	Jawa	2020)
personal		Tengah	
hygiene and			
pediculosis in Pondok			
Pesantren Al			
Yaqin			
Rembang			
Relationship	63.4% on	Karangan	(Pramadena,
of	santriwat	yar, Jawa	et al. 2020)
knowledge	i subjects	Tengah	,
level and	,	Ü	
personal			
hygiene to			
the			
incidence of			
pediculosis			
kapitis in boarding			
school			
Relationship	88.6% in	Palangka	(Rahmawati
of	orphana	Raya,	RK, et al.
knowledge	ge	Kalimanta	2020)
level and	children	n Tengah	,
behavior of		_	
shampoo			
use to the			
incidence of			
pediculosis			
kapitis in orphanage X			
Palangka			
Raya			
Relationship	60% in	Kota	(Maharani A,
of head	foster	Manado,	et al. 2020)
hygiene with	children	Sulawesi	,
pediculosis	and	Utara	
kapitis in the	street		
wall	children		
community in Bersehati			
Manado			
market			
Factors that	64.5% in	Kec.	(Nurdiani CU,
affect	santri	Cipayung,	2020)
pediculosis	and	Jakarta	,
capitis in	populatio	Timur	
children in	n		
Pondok			
Pesantren			
Sirojan			

Mustaqim			
and Residents			
rw 3 Kel.			
Pondok			
Ranggon			
Effectivenes	87.8% on	Mempawa	(Islami AC, et
s of	the	h Timur,	al, 2020)
counseling	subject of	Kalimanta	
on personal	santriwat	n Barat	
hygiene and	I		
incidence of			
pediculosis			
capitis in			
daughters of			
MTs in			
Pondok			
Pesantren X	44.55		
The	44.3% on	Jember,	(Nadira WA,
relationship between	populatio	Jawa Timur	et al. 2020)
personal	n subjects	Hilliui	
hygiene and	Jubjecto		
occupancy			
density with			
the			
incidence of			
pediculosis			
kapitis in Sukogidri			
Village			
Prevalence	78.5% on	Langowan	(Massie MA,
of pediculus	elementa	Timur,	et al. 2020)
humanus	ry school	Sulawesi	
capitis in	subjects	Utara	
elementary school			
children in			
East Kec.			
Langowan			
Relationship	38.3% on	Bantul,	(Sulistyani N,
of	elementa	Yogyakart	et al. 2019)
pediculosis	ry school	а	
kapitis,	subjects		
anemia status, and			
status, and learning			
achievement			
in			
elementary			
school			
students			
The	79.4% on	Bengkulu,	(Sudarsono,
relationship	santri	Sumatera	et al. 2019)
between	subjects		

personal		
hygiene and		
the		
incidence of		
pediculosis		
capitis in		
santriwati		
pondok		
pesantren		
Pancasila		
Bengkulu		

DISCUSSION

Research on pediculosis capitis is mostly done in Islamic boarding schools because of the density of residents and close contact. Based on these studies, it can be seen that the incidence of Pediculosis capitis in dormitories and orphanages is very high. This is due to the fact that Pediculosis capitis mainly attacks young children and guickly spreads in crowded living environments, for example in dormitories and orphanages. Based on the review that the author did, there are several things that make the incidence of Pediculosis capitis in Islamic boarding schools very high, namely direct transmission of Pediculosis capitis is very easy because female students often sleep together. In addition, there are several things that facilitate indirect transmission where female students in Islamic boarding schools often wear combs, towels and clothes together, borrow headscarves and mukena, this is what causes the incidence of Pediculosis capitis to be very high.3

The high incidence of *P. humanus capitis* infestation in girls may be due to the fact that girls have longer hair. Based on the observations in this review that most of the female students have longer hair. This allows transmission/transfer of the head tumor of Pediculus humanus capitis from one child to another. In line with other researchers, the frequency of Pediculosis capitis is more common in women than men, because women generally have longer hair than men. In addition, girls have a habit of playing with their friends in small groups and more often experience closer contact with their friends. The prevalence of infestations can vary depending on the policy in handling the incidence of infestation, the eradication method used, the number of direct contacts (head to head contact), the level of personal hygiene, living conditions and family economic conditions including family income as well as knowledge and attention in eradicating head lice.4

Pediculosis capitis causes the main clinical symptom in the form of itching on the scalp. The itching is caused by the injection of the tick's saliva into the scalp and causes an allergic reaction. Excessive itching causes sleep disturbances due to increased tick activity at night.⁵ Persistent sleep disturbances will have a negative impact in the form of decreased concentration power, decreased sharpness of memory, sensory, motor, and cognitive.⁶ Intense itching causes sleep disturbances and causes impaired concentration so that the patient's performance in daily activities is disrupted. Intense itching can disrupt sleep and interfere with learning concentration so that children's achievement decreases.⁷

Pediculosis capitis in addition to causing abnormalities in the hair and scalp can also cause other health problems in the form of iron deficiency anemia. Severe infections of the scalp caused by pediculosis capitis can cause a local inflammatory reaction in the form of regional lymph node enlargement (lymphadenopathy), especially in the occiput and retroauricular.⁸ Severe infestations of pediculosis capitis cause complications in the form of iron deficiency anemia. Iron deficiency anemia which is a complication of pediculosis capitis causes patients to feel lethargic, sleepy in class and affects their learning performance and cognitive function which has an impact on their learning achievement.⁹

REFERENCES

- Suweta NPTB, et al. PREVALENSI PEDICULOSIS CAPITIS DAN FAKTOR RISIKO INFESTASINYA di SD No 6 Darmasaba pada April – Mei 2018 . 2021;10(6):54–60.
- Fikri M, Siregar SM. JURNAL ILMIAH KOHESI Vol. 4 No. 2 April 2020. J Ilm Kohesi. 2020;4(2):100–5.
- 3. Anshor P, Analdi V, Santoso ID. Pediculus humanus capitis. Med Biol Illus. 2020;25(1):28.
- 4. Sulistyaningtyas AR, Ariyadi T, Zahro' F. Hubungan Antara Personal Hygiene dengan Angka Kejadian Pediculosis di Pondok Pesantrean Al Yaqin Rembang. 2020;9(1):25–31.
- 5. Sri K, Basuki W. HUBUNGAN TINGKAT PENGETAHUAN DAN PERSONAL HYGIENE

- TERHADAP INSIDEN PEDIKULOSIS KAPITIS DI The Relationship Level Knowledge and Personal Hygiene Against Pediculosis. 2019;016:326–34.
- 6. Rahmawati RK, Teresa A, Mutiasari D, Jelita H, Augustina I. Hubungan Tingkat Pengetahuan dan Perilaku Penggunaan Sampo Terhadap Kejadian Pedikulosis Kapitis di Panti Asuhan X Palangka Raya. J Kedokt. 2020;8(1):2–8.
- 7. Maharani A, Pandaleke HEJ, Niode NJ. Hubungan Kebersihan Kepala dengan Pedikulosis Kapitis pada Komunitas Dinding di Pasar Bersehati Manado. e-CliniC. 2019;8(1):163–71.
- 8. Nurdiani CU. Faktor-Faktor Yang Mempengaruhi Pediculosis Capitis Pada Anak-Anak Umur 6-12 Tahun Di Pondok Pesantren Sirojan Mustaqim Dan Penduduk Rw 03 Kelurahan Pondok Ranggon Kecamatan Cipayung Jakarta Timur. Anakes J Ilm Anal Kesehat. 2020;6(1):39–48.
- 9. Islami AC, Natalia D, Zakiah M. Efektifitas Penyuluhan Menggunakan Media Audia Visual Terhadap Personal Hygine dan Angka Kejadian Pediculosis Capitis pada Santri Putri Madrasah Tsanawiyah MTSn di Pondok Pesantren X Kecamatan Mempawa Timur. J Nas Ilmu Kesehat (Jnik) [Internet]. 2020;1(69):1–16. Available from: http://journal.unhas.ac.id/index.php/jnik/issue/view/542
- Nadira WA, Sulistyaningsih E, Rachmawati DA. Hubungan antara Personal hygiene dan Kepadatan Hunian dengan Kejadian Pedikulosis Correlation between Personal hygiene and Household Overcrowding to the Incidence of. J agromedicine Med Sci. 2020;6(3):161–7.
- Massie MA, Wahongan GJP, Pijoh V. Prevalensi Infestasi Pediculus humanus capitis pada Anak Sekolah Dasar di Kecamatan Langowan Timur. J Biomedik Jbm. 2019;12(1):24–30.
- 12. Sudarsono, S., & Miguna S. Hubungan antara personal hygiene dengan angka kejadian pediculosis capitis pada santriwati Pondok Pesantren Pancasila Bengkulun tahun 2018. J Univ Batam. 2019;9(1):81–91.